

- 堤 裕昭 (2005) 干潟の底質環境の変化とベントス群集への影響—有明海の砂質干潟を例として—. 月刊海洋 37: 107-115.
- 堤 裕昭・木村千寿子・永田紗矢香・佃 政則・山口一岩・高橋 徹・門谷 茂 (2004) : 広域定期観測による有明海水環境の現状, 沿岸海洋研究, 42: 35-42.
- Yamaguchi, H., Tsutsumi, H., Tsukuda, M., Nagata, S., Kimura, C., Yoshioka, M., Shibanuma, S. and S. Montani (2004). Utilization of photosynthetically produced organic particles by dense patches of suspension feeding bivalves on the sand flat of Midori River estuary, Kyushu, Japan. Benthos Research 59: 67-77.
- 堤 裕昭・岡村絵美子・小川満代・高橋徹・山口一岩・門谷 茂・小橋乃子・安達貴浩・小松利光 (2003) 有明海奥部海域における近年の貧酸素水塊および赤潮発生と海洋構造の関係. 海の研究 12: 291-305.
- Wu, S. S., Tsutsumi, H., Kita-Tsukamoto, K., Kogure, K., Ohwada, K., Wada, M. (2003) Visualization of the respiring bacteria in sediments inhabited by *Capitella* sp. 1. Fisheries Science 69: 170-175.
- Yamaguchi, H., Montani, S., Tsutsumi, H., Hamada, K., Ueda, N. (2003). Estimation of particulate organic carbon flux in relation to photosynthetic production in a shallow coastal area in the Seto Inland Sea. Mar. Pollut. Bull. 47: 18-24.
- Tsutsumi, H., Tsukuda, M., Yoshioka, M., Koga, M., Shinohara, R., Choi, K.-S., Cho, S.- C., Hong, J.-S. (2003) Heavy metal contamination in the sediment and its effect on the occurrence of the most dominant bivalve, *Ruditapes philippinarum* on the tidal flats of Ariake Bay in Kumamoto prefecture, the west coast of Kyushu, Japan. Benthos Research 58: 121-130.
- 堤 裕昭、石澤紅子、富重美穂、森山みどり、坂元香織、門谷 茂 (2002) 熊本有明地区の緑川河口干潟における盛砂後のアサリ (*Ruditapes philippinarum*) の個体群動態。日本ベントス学会誌 57: 177-187.
- 門谷 茂・山口一岩・堤 裕昭・濱田健一郎・上田直子 (2002) 瀬戸内海・周防灘における夏季リンの分布と生物生産環境. 沿岸海洋研究 39: 163-171.
- Tsutsumi, H., Montani, S., Kobe, H. (2002). Bioremediation of organic matter loaded on the sediment in outdoor pools with a polychaete, *Capitella* sp. 1. Fisheries Science 68: 613-616.
- Tsutsumi, H., Wainright, S., Montani, S., Saga, M., Ichihara, S., Kogure, K. (2001) Exploitation of a chemosynthetic food resource by a polychaete, *Capitella* sp. 1. Mar. Ecol. Prog. Ser. 216: 119-127.
- Tsutsumi, H., Kono, M., Hozumi, T., Hirashima, S., Takai, K., Manabe, T., Haraguchi, M., Yamamoto, I., Oppenheimer, C. (2001) Field tests of bioremediation agent with microbiological cultures. In: In Situ Aeration and Aerobic Remediation 6(10), ed. by Leeson, A., Johnson, P. C., Hinchee, R. E., Semprini, L., Magar, V. S., Battelle Press, pp. 345-352.
- Maguni, P., Montani, S., Takada, C., Tsutsumi, H. (2000). Extrapolation of

laboratory experiments on bivalve nutrient excretion to a field community situation: temporal scaling and relevance of biogenic nutrient flux from the sediments on a tidal flat of the Seto Inland Sea, Japan. *Mar. Ecol. Prog. Ser.* 198: 139-155.

Hozumi, T., Tsutsumi, H., Kono, M. (2000). Bioremediation on the shore after an oil spill from the *Nakhodka* in the Sea of Japan. I. Chemistry and characteristics of the heavy oil loaded on the *Nakhodka* and biodegradation tests on the oil by a bioremediation agent with microbiological cultures in the laboratory. *Mar. Pollut. Bull.* 40: 308-314.

Tsutsumi, H., Hirota, Y., Hirashima, A. (2000). Bioremediation on the shore after an oil spill from the *Nakhodka* in the Sea of Japan. II. Toxicity of a bioremediation agent with microbiological cultures in aquatic organisms. *Mar. Pollut. Bull.* 40: 315-319.

Tsutsumi, H., Kono, M., Takai, K., Manabe, T., Haraguchi, M., Yamamoto, I., Oppenheimer, C. (2000) Bioremediation on the shore after an oil spill from the *Nakhodka* in the Sea of Japan. III. Field tests of a bioremediation agent with microbiological cultures for the treatment of an oil spill. *Mar. Pollut. Bull.* 40: 320-324.

Ueda, N., Tsutsumi, H., Yamada, M., Hanamoto, K., Montani, S. (2000) Impact of oxygen-deficient water on the macrobenthic animals in the innermost areas of Dokai Bay and on the mud flats adjacent to the bay, in Kitakyushu, Japan. *Mar. Pollut. Bull.* 40: 906-913.

堤 裕昭、竹口知江、丸山 渉、中原康智 (2000) アサリの生産量が激減した後の緑川河口干潟に生息する底生生物群集の季節変化。日本ベントス学会誌 54: 1-8.

堤 裕昭 (2000) 1.2 九州の干潟と生物。In: 日本の水環境。7. 九州・沖縄編、日本水環境学会編、技報堂出版、東京、p. 9-14.

上田直子、堤 裕昭、明石久美、林田千恵美、山田真知子 (1999) 紫川河口域における底生動物の分布とその季節変化に影響を及ぼす環境要因について。日本ベントス学会誌 54: 4-13.

堤 裕昭 (1998) イトゴカイで海底の有機汚泥を浄化する。アクアネット 1: 50-53.

堤 裕昭、濱田建一郎、上田直子、山田真知子、藤木智子、中村仁美、和田育子、徳田貴子、門谷 茂 (1998) 富栄養化した水域の生態学的環境修復。—北九州市洞海湾を例として— *Capitella* sp. 1 (イトゴカイ、多毛類) を用いた洞海湾湾奥部の底質環境の修復。環境科学会誌 11: 421-429.

堤 裕昭・井上貴子 (1996) 20 年間にわたって大量の有機物負荷がかかってきた内湾域の海底環境と底生生物群集の特徴。日本ベントス学会誌, 50: 39-49.

門谷 茂・堤 裕昭 (1996) ベントスによる漁場底泥の環境修復。In: 生物機能による環境修復、石田祐三郎・日野明德編、恒星社厚生閣、東京、pp. 65-78.

Tsutsumi, H. (1995). Environmental impact of fish net pen culture on bottom of a cove, South Japan. *Estuaries* 18: 108-115.

- Chareonpanich, C., Tsutsumi, H., Montani, S. (1994). Efficiency of the decomposition of organic matter, loaded on the sediment, as a result of the biological activity of *Capitella* sp. I. Mar. Pollut. Bull. 28: 314-318.
- Chareonpanich, C., Montani, S., Tsutsumi, H., Nakamura, H. (1994). Estimation of oxygen consumption of a deposit-feeding polychaete *Capitella* sp. I. Fisheries Science 60: 249-251.
- Ueda, N., Tsutsumi, H., Yamada M., Takeuchi, R., Kido, K. (1994). Recovery of the marine bottom environment of a Japanese bay. Mar. Pollut. Bull. 28: 676-682.
- Chareonpanich, C., Montani, S., Tsutsumi (1994). Roles of a deposit-feeding polychaete, *Capitella* sp. I on the biological and chemical changes of the experimental marine sediment systems. Technical Bull. Fac. Agri. Kagawa Univ. 46: 21-26.
- Chareonpanich, C., Montani, S., Tsutsumi, H., Nakamura, H. (1994). An attempt to use a deposit feeding polychaete, *Capitella* sp. I, as a purifier of the bottom sediments in coastal fishery areas. Ed. by Chou, L. M., Munro, A. D., Lam, T. J., Chen T. W., Cheong, L. K. K., Ding, J. K., Hooi, K. K., Khoo, H. W., Phang, V. P. E., Shim, K. F., Tam, C. H. The Third Asian Fishery Forum, 31-34.
- Chareonpanich, C., Montani, S., Tsutsumi, H. (1993). Modification of chemical characteristics of organically enriched sediment by *Capitella* sp. I. Mar. Pollut. Bull. 26: 375-379.
- 堤 裕昭・門谷 茂 (1993) 魚類養殖場直下に堆積したヘドロ (有機汚泥) に対する生物浄化の試み。日本水産学会誌, 59: 1343-1347.
- 上田直子・堤 裕昭・徳崎健史・竹内良治・城戸浩三 (1992). 洞海湾湾岸部の底質環境と底生生物の分布。日本ベントス学会誌, 42: 55-62.
- Tsutsumi, H., Kikuchi, T., Tanaka, M., Higashi, T., Imasaka, K., Miyazaki, M. (1991). Benthic faunal succession in a cove organically polluted by fish farming. The Proceedings of International Conference on Environmental Management of Enclosed Coastal Seas. Mar. Pollut. Bull. 23: 233-238.
- Tsutsumi, H. (1990). Population persistence of *Capitella* sp. in a mud flat with environmental disturbance by organic enrichment. Mar. Ecol. Prog. Ser. 63: 147-156.
- Tsutsumi, H., Fukunaga, S., Fujita, N., Sumida, M. (1990). Relationship between growth of *Capitella* sp. and organic enrichment of the sediment. Mar. Ecol. Prog. Ser. 63: 157-162.
- Tsutsumi, H. (1987). Population dynamics of *Capitella capitata* (Polychaeta; Capitellidae) in an organically polluted cove. Mar. Ecol. Prog. Ser. 36: 139.
- 堤 裕昭 (1987). イトゴカイ (*Capitella capitata*) の有機物汚染域への適応。海洋科学, 19: 106-111.
- Tsutsumi, H., Kikuchi, T. (1984). Study of the life history of *Capitella capitata* (Polychaeta: Capitellidae) in Amakusa, South Japan including a comparison

with other geographical regions. Mar. Biol. 80: 315-3